

O² | âµ | · \$*!+ · ©, ©· wº !wu!

film©l :: » — ☰ — fl

0° — □ — — q^{®3} TM ± Ø — — — œº — — □ mœ³² Ñ ® | — — — a ² — μw ± Ø — ° μ ® — ³ | © » Q

q®³™

$q^{\text{R3-TM}} = \mu w - \ll$

» © nœc - R⁰ — R³ | . . R⁰ o - ceu - mu - R⁰ — R³ Ž à x' 2 q - x | . R³ — ± 3 TM - 3 | . o - c » Q

3 TM © { - μw⁰ . - 3 }

1! - Š° 2 Ž- Š° 2 - μ ° ©† | ® { - œ° « ° Q

2!^2 »Ž n - Ž ñ œ'—° ú®—° ° œ³ Q

4! Y|—·—° « « ¹ © —® E | —·—·— m™ — mcY · ©' » n wQ

5! Ÿ È—. · —° 2 3 · - ® - Ź - - Ź ® - · | · - - Ÿ œ° - - » œ » wQ

$$\frac{1}{\mu^{1/4}}$$

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

• 'â©Ž 2 3 x -3 Ž 0 0 · μ -· 2 · -¥- pŽ -®®Ž 2 0 ® , 3 · -· w¥-· · o -© · » Q

a 2 — œ — ñ°

(1fl³ « -TM μ | i³ @TM μ — @YTM \times — q μ x — wQ x » × i² @^{C°} « » | ° @^{C°} q μ x — —μ @^a | o — — » | wQ
 3 « 2 | ° » | » @^{C°} q μ x — — 3 Q

(2fl³ « —®|| | | n œ « 1© » | w © « 1© —®| — q© « 1© n œ — — 2 2 0 ° — | 1® — 0 — | 1® — 0 — |

(4fl³ « · · · · · · mō« · · TMx · » · © · · TM · — · , « · » · · 2 · μ · i · 3 Q

 ® · , â ©©

© 2003 Microsoft Corporation. All rights reserved. Microsoft, the Microsoft logo, and "Windows" are either registered trademarks or trademarks of Microsoft Corporation in the United States and/or other countries.

$$\frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}}$$

$\rightarrow p \cdot \cdot \cdot , \alpha \cdot \cdot 1 - , \rightarrow Q \mu \frac{1}{4} \cdot \cdot \cdot \circ \cdot \cdot \cdot \text{op } i \cdot \cdot \cdot Q$

μ | Ø — — ®

1! - □ - - 2 | ⊕ { - 3 q⁸³ TM — ⊙ | - - ⊙ y³ » - | Ø⁻³ ® - | ⊙ | - | - n ° « ' · ' « | 2 - 2 TM | « - oe » wQ

$2! \cdot 3^{\text{TM}} - 3^{\text{TM}3} + |\theta^0| - \langle \circ^0 | \cdot 3^{\text{TM}} - \mu^{3^{\circ}} \rangle \cdot \mu^{\circ} - \tilde{Y}|Q$

©

2! i ° -2 .. 3 n μ² .. 2 .. « , — ± Ø; .. — » m c Y ° .. n Z ° .. Z .. , ^ 3 a C ° .. « .. C a .. B b C μ u ° .. œ .. œ .. œ » w Q i ° .. — .. « , — ± Ø .. « .. 1 .. » - — .. - » Q

'o 3 —'o 'oe - -

- 2 © .

$m_j \cdot 2 \cdot \mu^- \cdot TMB \cdot m_0$

qñ® « - ® 'o ' -

$m_i \cdot 2 \cdot \mu^{-1} \cdot TMB \cdot m_o$

Caution:† Not to Be Used on Crops Other Than Specified on This Label/Leaflet†

IMIDACLOPRID 17.8% SL**(Systemic Insecticide)**

Imidacloprid 17.8% SL is a systemic insecticide containing Imidacloprid active ingredient balance auxiliaries and inert material. It is used as a spray for the control of sucking and other insects in different crops.

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water (Ltr)	Waiting Period between last spray to harvest (in days)
		AI (kg)	Formulation (Ltr)		
Grapes	Flea beetle	0.06-0.08	0.300-0.400	1000	32
Mango	Hopper	0.4-0.8g/tree	2-4g/tree	10	45
Cotton	Aphid, whitefly, Jassids, Thrips	0.020-0.025	0.10-0.125	500-700	40
Sugarcane	Termite	0.07	0.35	1875	45
Tomato	White fly	0.03-0.035	0.150-0.175	500	3
Paddy	BPH, WBPH, GLH	0.020-0.025	0.10-0.125	500-700	40
Citrus	Leaf miner, psylla	0.01%	0.05%		15
Okra	Aphid, Jassids, Thrips	0.02	0.1	500	3
Chillies	Jassid, Aphid, Thrips	0.025-0.05	0.125-0.250	500-700	40
Groundnut	Aphid, jassids	0.02-0.025	0.1-0.125	500	40
Sunflower	Jassids, Thrips, whitefly	0.02	0.1	500	30

Direction of Use

Plant Protection Equipment: Hand operated knapsack sprayer or foot sprayer is recommended and tractor mounted sprayer can also be used.

Precaution

1. Keep away from foodstuffs, empty foodstuff containers and animals feed.
2. Avoid contact with eyes, mouth and skin.
3. Avoid inhalation the spray mist. Spray in the direction of wind.
4. Wash thoroughly the contaminated clothes and part of the body after spraying.
5. Do not smoke, drink eat and chew anything while spraying.
6. Wear full protective clothing while mixing and spraying.

Symptoms Of Poisoning

Apathy, myotonia, tremor, difficult breathing, myospasms may occur.

First Aid

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet ..

1. If swallowed, induce vomiting by tickling the back of throat. Repeat it until the vomitus is clear. Do not induce vomiting if the patient unconscious.
2. If clothing and skin is contaminated, remove the clothes and wash the contaminated skin with copious amount of soap and water.
3. If eyes are contaminated, flush with plenty of saline/clean water for about 10 to 15 minutes.
4. If inhaled, remove the patient to fresh air.

Phytotoxicity

The product is not Phytotoxic when used as per field recommendation.

Antidote

No specific antidote. Treat symptomatically.

Disposal Of Used Container

1. The packages shall be broken and buried away from habitation.
2. The used packages shall not be left outside to prevent their re-use.
3. Packages or surplus material and washing should be disposed in a safe manner so as to prevent environment or water pollution.

Storage Conditions

- 1) The package containing the insecticide shall be stored in separate rooms or premises away from the rooms or premises used for stored other articles shall be kept in separate Almirahs under lock and key, depending on the quantity and nature of insecticides
- 2) The rooms or premises meant for storing the insecticide shall be well built, dry, and well lit and ventilated and sufficient dimensions to avoid contamination with vapour.

Chemical Composition:

Imidacloprid	(a.i.)	17.80 % w/w
Non ionic emulsifier	(Ethoxylated Alkyl Aryl Phenol Derivative)	2.50 % w/w
Polyvinyl Pyrrolidone Copolymer		1.00 % w/w
Dimethyl Sulfoxide		38.40 % w/w
N-Methyl Pyrrolidone		Q.S % w/w
Total:		100.000% w/w

Manufactured By:

Toshi Insecticides India
Gaushala Road, Near Cooler Factory, Geeta House, Karnal
Karnal, Karnal, Haryana

Manufacturer Premises :

Meerut Road, Village Andhera, District-Karnal, Haryana-132001

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet ..