

$\text{fl} = \varnothing^3 = \circ \mid \text{fl} \neq !\text{w}^2 \neq \text{!fl}$ fl \rightarrow μ \rightarrow fl

$\circ \text{!w}^2 = \text{fl} = \circ \text{!fl}$ ($\# \text{!} \circ \text{!w}^2 \text{!} \text{fl} \text{!} \text{fl}$) $\circ \text{!} \text{fl} = (\# \text{!} \circ \text{!w}^2 \text{!} \text{fl})$

$\text{!} \text{w}^2 = \text{fl} = \text{!} \text{fl}$

$\text{!} \text{!w}^2 = \text{fl} = \text{!} \text{fl}$

$\text{!} \text{w}^2 = \text{fl} = \text{!} \text{fl}$

$\text{!} \text{w}^2 = \text{fl} = \text{!} \text{fl}$

$\text{!} \text{w}^2 = \text{fl} = \text{!} \text{fl}$

q^{®3} TM

μ	$\text{fl} = \mu \text{!} \text{fl}$			
μ	$\text{fl} = \mu \text{!} \text{fl}$			
μ	$\text{fl} = \mu \text{!} \text{fl}$			

q^{®3} TM — $\mu \text{!} \text{fl}$ — Q

$\text{!} \text{!} \text{w}^2 = \text{fl} = \text{!} \text{!} \text{w}^2$

$\text{!} \text{!} \text{w}^2 = \text{fl} = \text{!} \text{!} \text{w}^2$

$\text{!} \text{!} \text{w}^2 = \text{fl} = \text{!} \text{!} \text{w}^2$

$\text{!} \text{!} \text{w}^2 = \text{fl} = \text{!} \text{!} \text{w}^2$

$\text{!} \text{!} \text{w}^2 = \text{fl} = \text{!} \text{!} \text{w}^2$

$\text{!} \text{!} \text{w}^2 = \text{fl} = \text{!} \text{!} \text{w}^2$

3 TM — $\text{fl} = \mu \text{!} \text{fl}$ — Q

$\text{!} \text{!} \text{w}^2 = \text{fl} = \mu \text{!} \text{fl}$

$\text{!} \text{!} \text{w}^2 = \text{fl} = \mu \text{!} \text{fl}$

$\text{!} \text{!} \text{w}^2 = \text{fl} = \mu \text{!} \text{fl}$

$\text{!} \text{!} \text{w}^2 = \text{fl} = \mu \text{!} \text{fl}$

$\text{!} \text{!} \text{w}^2 = \text{fl} = \mu \text{!} \text{fl}$

$\text{!} \text{!} \text{w}^2 = \text{fl} = \mu \text{!} \text{fl}$

$\text{!} \text{!} \text{w}^2 = \text{fl} = \mu \text{!} \text{fl}$

. $\text{fl} = \mu \frac{1}{4} \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

a 2 — oe-ñ⁰

$\text{fl} = \mu \text{!} \text{fl}$

. $\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

$\text{fl} = \mu \text{!} \text{fl}$

. $\mu \text{!} \text{fl}$ **Caution:**“Not to Be Used on Crops Other Than Specified on This Label/Leaflet”.

1! " μ | Ø | Ø | — ⊙ | — qō» n ° « '° ' « ' i 2 - ' 2 ' « ' - ' « Q
2! " μ | Ø | Ø | — @— 3 TM | — μ w qō» ° » ' - ' — Q
3! " μ | Ø | Ø | 3 ° œ » n ° 2 - ' Ø a ' 2 , - ' E Ø | — ' — ' — Ø a ' 0 . n - @ | — ' — Ø 3 | ' — ' Ø a ' i μ ' « 1 " - ' » Q

° » — ⊙

1! " — Ø | Ø | — ' 0 » ' mμ TM — 2 ' 3 ' a - ' Ø mō 3 ' | ©{ ' ' , 1 — ' — ' — Ø a ' — ' 0 » ' — 1/4 ' 0 ' « ' — 3 ' i - ' œ » w 3 ' — Ø | — ' 2 ' | ' — ' ± ' — Ø ' — ± ' — ' © » w E Ø | — μ » ' mμ 2 ' — ' 2 ' © μ Žœ ' ' μ TM — ' — Q ' 2! " — Ø | — ' 0 » ' — 1/4 ' 3 ' a - ' , - « ' Ž 2 ' i ° © Ž » ' « ' © a ' — Ø 3 | ' μ ú ' p Žœ E p | — ' œ » w Q — 1/4 — ' a © ' — ' ¥ E ' » - ' œ » w Q

° 0 3 — ' 0 ' œ — -

% Z° — Ž 3 Ø Ø | — ' 3 μ ' w — Ž 2 a Ø μ ' (#!# # ' ' ± ' " ± ' — ' P Ø Ž° @ w 2 ' 0 ' ! ' © ! ' f Ø 3 — Ø 2 ' , © ' , (' © , © fl
0² μ ' 3 ' — Ø Ø | — ' — y , 3 2 ' w y — Ø μ ' \$#!# # ' ' ± ' " ± ' w ' 3 μ ' 0 y - — Ø w Ø | — Ø μ ' n a ' o a μ - ' p a ' ° μ ' Ø w ' 2 a ' fl © 3 | — 2 ' ' — ' ± ' " ± '

— 2 Ø |

m | — ' 2 ' μ — ' TM 3 | m — ' —

gñ@ « - ' @ ' 0 ' -

m | — ' 2 ' μ — ' TM 3 | m — ' —

Caution:“Not to Be Used on Crops Other Than Specified on This Label/Leaflet”.

Fenobucarb (B.P.M.C) 50% EC**(Insecticide)**

BPMC (Fenobucarb) 50 % E.C is a stomach and contact action insecticides recommended for the control of Brown Plant hopper, Green leaf hopper of rice.

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water (Litr)	Waiting Period between last spray to harvest (in days)
		AI (Kg)	Formulation (ltr)		
Paddy	Brown Plant Hopper, Green Leaf Hopper	0.25-0.75	0.5-1.5	500	30

Direction of Use

Knapsack sprayer, Foot Sprayer, Compression Knapsack sprayer Compression knapsack, battery sprayer and ASPEE-HTP power sprayer.

Precaution

1. Do not mix the pesticides with bare hands.
2. The user should use full protective clothing which includes rubber gloves, boots either face shield or dust mask and an over all rubber apron hood or hat.
3. Use in high concentration with low and volume, application equipment is dangerous and should be avoided.

Symptoms Of Poisoning

Salivation, watery eyes, pinpoint eye pupils, blurred vision, muscles tremors, difficulty in breathing, excessive sweating, abdominal cramps, nausea, vomiting diarrhea, weakness, headache. In severe cases, convulsions, unconsciousness and respiratory failures may occur.

First Aid

In case of poisoning, the patient should be removed to fresh air. If swallowed, induce vomiting by tickling the back of throat. If skin or clothes are contaminated. Remove the contaminated clothes immediately and wash affected parts with soap and water. If eyes are affected wash with water. In case of accidental poisoning call a physician.

Antidote

Atropine sulphate administers Intramuscularly or orally .The patient should be observed for several hours following exposures and atropine sulphate may be re- administered if symptoms persist or occur.

Disposal Of Used Container

1. The packages shall be broken and buried away from habitation.
2. The packages shall be not left to prevent their re-use.
3. The used packages or surplus materials and washing from the machine the container shall be disposed off in safe manner so as to prevent environmental and water pollution.

Storage Conditions

1. The packages containing the insecticide, shall be stored in separate rooms or premises away from the rooms or premises used for storing other articals or shall be kept in safe manner in almirahs under lock and key depending upon the quantity and nature of the insecticides.
2. The rooms or premises meant for storing the insecticide shall well built, dry, well-lit and ventilated and of sufficient dimensions to avoid contaminating with vapours.

Caution:“Not to Be Used on Crops Other Than Specified on This Label/Leaflet”.

Chemical Composition:

2-Sec-Butyl phenyl N-methyl carbamate (BPMC) a.i. (Min. purity 95%)	50.00 % w/w
Emulsifiers : Blend of Calcium alkyl aryl sulphonate & Poly oxy ethylene ether	10.00 % w/w
Solvent (Aromax)	QS % w/w
Total:	100.000% w/w

Manufactured By:

Toshi Insecticides India
Gaushala Road, Near Cooler Factory, Geeta House, Karnal
Karnal, Karnal, Haryana

Manufacturer Premises :

Meerut Road, Village Andhera, District-Karnal, Haryana-132001

Caution:“Not to Be Used on Crops Other Than Specified on This Label/Leaflet”.