

— °ō! j 2 '\$% 'f 2 --j °) & 'šμ-, μ'@q! '!

fi m@l . » 'w ' ' @, 3 ' ' « - , — fl

— °ō! j 2 '\$% 'f 2 --j °) & 'šμ-, μ'@q! 'w— mñ³ ò@ ± , μ '¼— w 2 ° ò³ ' ' « - , — » Q³ »' ± , μ ' @ ' — ° ' — ' @ ' μ™— . μ ' ° » 2 ' 2™ μ ' @ ' μ™— . μ ' @ ' šø° ' . ° » 2 ' 2™ Ž n μ ' @ ' μ™— . μ ' mμ ' @μ ož μ ' @μ ož μ — ° — ' . ' œ³ ' @ ' μ™— . μ ' ' @μ ož ' @μ ož ' ' ' ž | p' ° — ž @μ — ' @ ' μ™— . — 3 ' ' — ' @ ' » Q

q@3™

° μ	— — 2	° » â ³ ' 2		@ — — 2 fμ ' fl	m @² ' Ÿ — . @ ' ° μ — — ° œ mō ' μ fi « — fl
		° ! @ñ fi 2 fl	° ' œ— fi 2 fl		
m™	2 « ' 2 @ ' n ° @ ' '™ž œ ' @ ' n ° @ ž wō â — j	0.11%	0.15%	' ° μ ' — — @ ' — m— ° ' n . 3 — 2	7
n 2	œ ' @ ' n ° @ ž wō â — j	0.11%	0.15%	' ° μ ' — — @ ' — m— ° ' n . 3 — 2	7
n μ	mμ ' @μ ož μ ' @μ ož μ — ° —	1312.5	1750	500	47
œ³	@μ ' @μ ož ' @μ ož ' ' ž p' ° — ž @μ — ' @	937.5	1250	250-500	7
— —	° » 2 ' °	563	750	750	57
2™ μ	@ ' šø° ' ž ° » 2 ' °	375	500	500	72

q@3™ — μ w — « —

@Ÿ ' @ ' μ « — . μ ' Ÿ | — . — — 3 ò ' @ ' qçœ n ³ @— . μ ' Ÿ | — . 3 ò ' — 3™ — Q

3™ — ° 2 3 ' °

o ' ' « - , — — 3™ '™ — μ¼ ' « — ' @ ' @ ' @ ' Ÿ | — . — Q — « , ' m— ° ' ' « - , — w 2 ' @ — — q œ @ 2 ' 2 ' 2 ' — — Ÿ | — . — Q ' « - , — — Ÿ | — . ° » μž o° — š μ — mçŸ ' @ ' w— μ — ' . ' 2 μ ' μ Q

3™ @ (— μ w ° . — 3 ' °

Caution:Í Not to Be Used on Crops Other Than Specified on This Label/LeafletÍ™

— ° ö! j 2 ° á — μ	, + ° ! ©!	\$%&(' ± " ± '
2 — j ° ° á — μ	+(' ° ! ©!	*' ! \$%(' ± " ± '
° 3 2 ° ý ž v μ — μ ' w o μ ' ° ý - — α		%##(' ± " ± '
° 3 2 ° ý ž v μ — μ ' - ® o μ ' ° ý - — α		%##(' ± " ± '
— n μ - ' - 3 ' ® α a		, !) &(' ± " ± '

3 TM - TSS!SSS(' ± " ± '

-2 ©

m j ' 2 ' μ - ' TM B ' m - ° '

qñ® « - ' ® ' ° ' -

m j ' 2 ' μ - ' TM B ' m - ° '

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Carbendazim 12% + Mancozeb 63% WP**(Systemic and Contact Fungicide)**

Carbendazim 12%+ Mancozeb 63% WP is very effective, crops protective and curative fungicides. It successfully control of blast disease of paddy, leaf spot and blast disease of groundnut, early blight, late blight, black scurf of potato, blister blight, grey blight, red rust, die back & black rot or tea crop.

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water (Litr)	Waiting Period between last spray to harvest (Days)
		AI (gm)	Formulation (gm)		
Grape	Downy mildew, powdery mildew, anthracnose	0.11%	0.15%	As required depending on crop canopy	7
Mango	powdery mildew, anthracnose	0.11%	0.15%	As required depending on crop canopy	7
Potato	Early blight, Late blight, Black scurf	1312.5	1750	500	47
Tea	Blister blight, Grey blight, Red rust, Die-back, Black rot	937.5	1250	250-500	7
Paddy	Blast	563	750	750	57
groundnut	leaf spot, blast	375	500	500	72

Direction of Use

Spray by using high volume sprayer viz. Kmapsack sprayer.

Time of Application

The product should be sprayed immediately on appearance of disease symptoms. As per recommendation, mix the pesticides and water at right dosage and spray. Before spraying the pesticides, It's suspension should be mixed well by a wooden stick.

Precaution

- 1) Keep away from foodstuffs, empty foodstuff containers and animals feed.
 - 2) Avoid contact with mouth, eyes and skin.
 - 3) Avoid inhalation of the spary mist. Spray in the direction of wind.
 - 4) Wash thoroughly the contaminated clothes and parts of the body after spraying.
 - 5) Do not smoke, drink eat and chew anything while spraying.
 - 6) Wear full protective clothing while mixing and spraying.
- CAUTION FOR WORKERS/USERS:- while using fungicide workers/users must wear full protective clothing like long sleeved shirt, long paint, chemical resistant gloves, shoes, goggles hat and mask.

Symptoms Of Poisoning

Sweating, headache, lethargy, coughing, sneezing, itching, nausea, vomiting may occur.

First Aid

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

- 1) If swallowed, induce vomiting by tickling the back of throat. Repeat it until the vomitus is clear. Do not induce vomiting if the patient is unconscious.
- 2) If clothing and skin are contaminated, remove the cloths and wash the contaminate skin with copious amount of soap and water.
- 3) If eyes are contaminated, flush with plenty of saline/clean water for about 10 to 15 minutes.
- 4) If inhaled, remove the patient to fresh air.

Phytotoxicity

The product is not phytotoxic when used as per field recommendation.

Antidote

No antidote is known. Treat symptomatically.

Disposal Of Used Container

- 1) The package containing the fungicide shall be stored in separate rooms or premises away from the rooms or premises used for storing articles or shall kept in separate almirahs under lock and key depending upon the quantity and nature of fungicide.
- 2) The rooms or premises meant for storing the fungicide shall be well built, dry, well lit and ventilated and sufficient dimension.

Storage Conditions

Packages or surplus material and washing from the machines and cointainer shall be disposed off in such a manner so as to prevent environmental or water pollution. The used packages should not be left outside to prevent their re-use. Packages should be burried away from habitation and source of water.

Chemical Composition:

Carbendazim Technical	(based on 98% w/w a.i.	12.25 % w/w
Mancozeb Technical	(based on 85% w/w a.i.)	74.12 % w/w
Sodium Salt of alkyl aryl sulfonate		2.00 % w/w
Sodium Salt of alkyl naphthyl sulfonate		2.00 % w/w
Keolin		9.63 % w/w
Total:		100.000% w/w

Manufactured By:

Toshi Insecticides India
Gaushala Road, Near Cooler Factory, Geeta House, Karnal

Karnal, Karnal, Haryana

Manufacturer Premises :

Meerut Road, Village Andhera, District-Karnal, Haryana-132001

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet